

AFTER SURGERY

- You will be given drops to take home. Please ensure you follow the instructions. Wash your hands before applying the drops. Do not let the tip of the bottle touch your eye. You may experience tearing, glare and light sensitivity following surgery. Sunglasses may help with these symptoms. You will be provided an eye shield to take home, please use it especially when sleeping.



Wash
hands



Correct
application



Protective
sunglasses



Eye shield
when
sleeping

- Avoid strenuous activities such as heavy lifting (1-2 weeks). Swimming is to be strictly avoided (1 month). If you have a gas bubble in your eye, air travel and going up-hill is prohibited until your doctor clears it.



Avoid
heavy lifting



Swimming
prohibited



Ask your
Doctor

CHECKLIST

Name:

MRN No.:

Surgeon:

Procedure:

Estimated Price:

Deposit:

Remarks:

Post-op Emergency Contact:

Patient Signature:

Contact the Post-Op Emergency number above if you experience any of the following symptoms after your procedure:

- Worsening pain / redness in the operated eye
- Severe / sudden pain
- Sudden loss of vision
- Nausea or vomiting after surgery
- Accidental injury to the operated eye



CONTACT US



03 2730 7666



oasiseye.my



linktr.ee/oasiseye



YOUR EYES AND YOU

Preparing for your
eye surgery at



OASISEYE
SPECIALISTS

PROTECTING VISION | RESTORING SIGHT





BEFORE SURGERY

- Know the procedure you will be having. Do not hesitate to ask your doctor if you are unsure about any details.
- You will be required to pay a deposit once a date has been fixed.
- The full amount for the estimated price of the procedure will need to be settled prior to surgery.
- In the event of postponement, cancellation or excess payment, the deposit / unutilised portion will be refunded to you.
- If absolutely necessary, any unforeseen usage of medication / equipment during surgery will be added to the final bill.



ON THE DAY OF SURGERY

- Please be on time for your appointment. Do inform us if you need to postpone for any reason.



- Continue with your regular medication except those indicated by the doctor, such as blood-thinners.




- Have a shower before surgery, taking care to clean your face and around your eyes and lashes.



- Do not use any alcohol-based products such as cosmetics, hair products, nail polish and perfume.



- If your surgery is under local anaesthesia, you can have a light meal prior to surgery.



- If your surgery is under general anaesthesia, you cannot eat or drink for 6 hours before surgery.



- Bring as few valuables as you can. There is a small locker available for your personal use.



- You cannot drive yourself home after surgery. Please ensure you have someone to take you home.



- Dilating drops may be applied to your eyes which will blur your vision for a few hours.



- For hygiene purposes, you will be required to change into an OT gown for surgery.

